

SIMPLY CONNECTED SYMPLECTIC 4-MANIFOLDS WITH $b_2^+ = 1$ AND $c_1^2 = 2$

JONGIL PARK

ABSTRACT. In this article we construct a new family of simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$ which are not diffeomorphic to rational surfaces by using rational blow-down technique. As a corollary, we conclude that a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$ admits an exotic smooth structure.

1. INTRODUCTION

One of the fundamental problems in the topology of 4-manifolds is to determine whether a given topological 4-manifold admits a smooth structure and, if it does, whether such a smooth structure is unique or not. Though the complete answer is far from reach, gauge theory makes us to answer partially these questions ([FS1], [FS2], [G], [Sz], [T]). But most known results are in the case of simply connected 4-manifolds with either $b_2^+ > 1$ odd or $b_2^+ = 1$ and $c_1^2 \leq 0$. In the case when $b_2^+ = 1$ and $c_1^2 > 0$, a theorem of D. Kotschick is the only known result that the Barlow surface is not diffeomorphic to $\mathbf{CP}^2 \# 8\overline{\mathbf{CP}}^2$ ([K1]). Since then, there was little progress on the problems.

In this paper we investigate exotic smooth structures on a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$. According to a convention, we say that a smooth 4-manifold admits an *exotic smooth structure* if it has more than one distinct smooth structure. One way to get an exotic smooth structure on a given smooth 4-manifold X is to construct a new smooth 4-manifold X' which is homeomorphic, but not diffeomorphic, to X . Hence the problem of finding exotic smooth structures on a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$ is equivalent to find a new family of simply connected smooth 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$. Note that the only known simply connected closed smooth 4-manifolds with $b_2^+ = 1$ and $c_1^2 \geq 2$ are rational surfaces such as $\mathbf{CP}^2$, $S^2 \times S^2$ and $\mathbf{CP}^2 \# n\overline{\mathbf{CP}}^2$ ($n \leq 7$). Despite the fact that it is no constraint on the existence of simply connected smooth 4-manifolds with $b_2^+ = 1$ and $c_1^2 \geq 2$ which are not rational surfaces, no such 4-manifolds have been known. Thus it has long been an interesting question to find such 4-manifolds (refer to Problem 4.45 in the Kirby list appeared in [Ka]). In Section 3 we construct a new family of simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$ which are homeomorphic, but not diffeomorphic, to rational surfaces by using rational blow-down technique introduced by R. Fintushel and R. Stern in [FS1]. As one of our main results, we get the following

Theorem 1.1. *There exists a simply connected symplectic 4-manifold with $b_2^+ = 1$ and $c_1^2 = 2$ which is homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$.*

Date: Revised: March 23, 2004.

2000 Mathematics Subject Classification. Primary 53D05, 14J26; Secondary 57R55, 57R57.

Key words and phrases. Exotic smooth structure, rational blow-down, rational surface, symplectic.

This work was supported by grant No. R14-2002-007-01002-0 from KOSEF.

Hence we conclude that a rational surface $\mathbf{C}P^2 \# 7\overline{\mathbf{C}P}^2$ admits an exotic smooth structure. Furthermore, by blowing up of a symplectic 4-manifold constructed in Theorem 1.1 above, we also conclude that a rational surface $\mathbf{C}P^2 \# 8\overline{\mathbf{C}P}^2$ admits at least three distinct smooth structures - an Einstein metric with positive scalar curvature, an Einstein metric with negative scalar curvature and no Einstein metric.

Acknowledgements. The author would like to thank Ronald Fintushel for his constant encouragement and guidance. The author is deeply indebted to him for sharing his ideas with the author while working on this problem. The author would also like to thank András Stipsicz for pointing out some critical errors in the earlier version of this article and an explanation about various types of singular fibers appeared in a rational surface $E(1)$. The author also thanks Denis Auroux for figuring out $\widehat{E}_6$ -fiber in $E(1)$ and Tian-Jun Li for an explanation about the symplectic 2-forms on a non-minimal rational surface. Finally he also thanks the department of Mathematics at Michigan State University for hospitality and support during his visit.

2. PRELIMINARIES

In this section we briefly review the Seiberg-Witten theory and a rational blow-down surgery which will be the main technical tools to get our results.

First we briefly introduce the Seiberg-Witten theory for smooth 4-manifolds. In particular, we pay attention to the Seiberg-Witten invariant of 4-manifolds with $b_2^+ = 1$ ([M] for details). Let X be a closed, oriented smooth 4-manifold with $b_2^+ > 0$ and a fixed metric g , and let L be a characteristic line bundle on X , i.e. $c_1(L)$ is an integral lift of $w_2(X)$. This determines a $Spin^c$ -structure on X which induces a complex spinor bundle $W \cong W^+ \oplus W^-$, where $W^\pm$ is the associated $U(2)$ -bundles on X such that $\det(W^\pm) \cong L$. Note that the Levi-Civita connection on TX together with a unitary connection A on L induces a connection $\nabla_A : \Gamma(W^+) \rightarrow \Gamma(T^*X \otimes W^+)$. This connection, followed by Clifford multiplication, induces a $Spin^c$ -Dirac operator $D_A : \Gamma(W^+) \rightarrow \Gamma(W^-)$. Then, for each self-dual 2-form $h \in \Omega_{+g}^2(X; \mathbf{R})$, the following pair of equations for a unitary connection A on L and a section Ψ of $\Gamma(W^+)$ are called the *perturbed Seiberg-Witten equations*:

$$(1) \quad (SW_{g,h}) \begin{cases} D_A \Psi &= 0 \\ F_A^{+g} &= i(\Psi \otimes \Psi^*)_0 + ih. \end{cases}$$

Here F_A^{+g} is the self-dual part of the curvature of A with respect to a metric g on X and $(\Psi \otimes \Psi^*)_0$ is the trace-free part of $(\Psi \otimes \Psi^*)$ which is interpreted as an endomorphism of W^+ . The gauge group $\mathcal{G} := \text{Aut}(L) \cong \text{Map}(X, S^1)$ acts on the space $\mathcal{A}_X(L) \times \Gamma(W^+)$ by

$$g \cdot (A, \Psi) = (g \circ A \circ g^{-1}, g \cdot \Psi)$$

Since the set of solutions is invariant under the action, it induces an orbit space, called the *Seiberg-Witten moduli space*, denoted by $M_{X,g,h}(L)$, whose formal dimension is

$$\dim M_{X,g,h}(L) = \frac{1}{4}(c_1(L)^2 - 3\sigma(X) - 2e(X))$$

where $\sigma(X)$ is the signature of X and $e(X)$ is the Euler characteristic of X . Note that if $b_2^+(X) > 0$ and $M_{X,g,h}(L) \neq \emptyset$, then for a generic self-dual 2-form h on X the moduli space $M_{X,g,h}(L)$ contains no reducible solutions, so that it is a compact, oriented, smooth

manifold of the given dimension.

Definition The *Seiberg-Witten invariant (for brevity, SW-invariant)* for a smooth 4-manifold X with $b_2^+ > 0$ is a function $SW_X : Spin^c(X) \rightarrow \mathbf{Z}$ defined by

$$(2) \quad SW_X(L) := \begin{cases} \langle \beta^{d_L}, [M_{X,g,h}] \rangle & \text{if } \dim M_{X,g,h}(L) := 2d_L \geq 0 \text{ and even} \\ 0 & \text{otherwise.} \end{cases}$$

Here β is a generator of $H^2(\mathcal{B}_X^*(L); \mathbf{Z})$ which is the first Chern class of the S^1 -bundle

$$\tilde{\mathcal{B}}_X^*(L) = \mathcal{A}_X(L) \times (\Gamma(W^+) - \{0\}) / Aut^0(L) \longrightarrow \mathcal{B}_X^*(L)$$

where $Aut^0(L)$ consists of gauge transformations which are the identity on the fiber of L over a fixed base point in X . Note that if $b_2^+(X) > 1$, the Seiberg-Witten invariant, denoted by $SW_X = \sum SW_X(L) \cdot e^{c_1(L)}$, is a diffeomorphism invariant, i.e. SW_X does not depend on the choice of a metric on X and a generic perturbation of Seiberg-Witten equations. Furthermore, only finitely many $Spin^c$ -structures on X have a non-zero Seiberg-Witten invariant. We say that the characteristic line bundle L , equivalently a cohomology class $c_1(L) \in H^2(X; \mathbf{Z})$, is a *SW-basic class* of X if $SW_X(L) \neq 0$.

When $b_2^+(X) = 1$, the SW-invariant $SW_X(L)$ defined in (2) above depends not only on a metric g but also on a self-dual 2-form h . Because of this fact, there are several types of Seiberg-Witten invariants for a smooth 4-manifold with $b_2^+ = 1$ depending on how to perturb the Seiberg-Witten equations. We introduce two types of SW-invariants and investigate how they are related. First we allow all metrics and self-dual 2-forms to perturb the Seiberg-Witten equations. Then the SW-invariant $SW_X(L)$ defined in (2) above has generically two values which are determined by the sign of $(2\pi c_1(L) + [h]) \cdot [\omega_g]$, where ω_g is a unique g -self-dual harmonic 2-form of norm one lying in the (preassigned) positive component of $H_{+g}^2(X; \mathbf{R})$. We denote the SW-invariant for a metric g and a generic self-dual 2-form h satisfying $(2\pi c_1(L) + [h]) \cdot [\omega_g] > 0$ by $SW_X^+(L)$ and denote the other one by $SW_X^-(L)$. Then the wall crossing formula tells us the relation between $SW_X^+(L)$ and $SW_X^-(L)$.

Theorem 2.1 (Wall crossing formula, [M]). *Suppose that X is a closed, oriented smooth 4-manifold with $b_1 = 0$ and $b_2^+ = 1$. Then for each characteristic line bundle L on X such that the formal dimension of the moduli space $M_{X,g,h}(L)$ is non-negative and even, say $2d_L$, we have*

$$SW_X^+(L) - SW_X^-(L) = -(-1)^{d_L}.$$

By the way, C. Taubes' result on the SW-invariant of a symplectic 4-manifold with $b_2^+ > 1$ can be easily extended to the $b_2^+ = 1$ case.

Theorem 2.2 ([T], [LL1]). *Suppose X is a closed symplectic 4-manifold with $b_2^+ = 1$ and a canonical class K_X . Then $SW_X^-(K_X) = \pm 1$.*

Second one may perturb the Seiberg-Witten equations by adding only a small generic self-dual 2-form $h \in \Omega_{+g}^2(X; \mathbf{R})$, so that one can define the SW-invariants as in (2) above. In this case we denote the SW-invariant for a metric g satisfying $(2\pi c_1(L)) \cdot [\omega_g] > 0$ by $SW_X^{\circ,+}(L)$ and we denote the other one by $SW_X^{\circ,-}(L)$. Note that $SW_X^{\circ,\pm}(L) = SW_X^{\pm}(L)$. But it sometimes happens that the sign of $(2\pi c_1(L)) \cdot [\omega_g]$ is the same for all generic metrics,

so that there exists only one SW-invariant obtained by a small generic perturbation of the Seiberg-Witten equations. In such a case we define the SW-invariant of L on X by

$$SW_X^\circ(L) := \begin{cases} SW_X^{\circ,+}(L) & \text{if } 2\pi c_1(L) \cdot [\omega_g] > 0 \\ SW_X^{\circ,-}(L) & \text{if } 2\pi c_1(L) \cdot [\omega_g] < 0. \end{cases}$$

If $SW_X^\circ(L) \neq 0$, we call the corresponding $c_1(L)$ (or L) a *SW-basic class* of X . Then the Seiberg-Witten invariant of X , denoted by $SW_X^\circ = \sum SW_X^\circ(L) \cdot e^{c_1(L)}$, will also be a diffeomorphism invariant. Furthermore we can extend many results obtained for smooth 4-manifolds with $b_2^+ > 1$ to this case. For example, we have

Theorem 2.3. *Let X be a simply connected closed smooth 4-manifold with $b_2^+ = 1$ and $b_2^- \leq 9$. Then*

- (i) *There are only finitely many characteristic line bundles L on X such that $SW_X^\circ(L) \neq 0$.*
- (ii) *If X admits a metric of positive scalar curvature, then the SW-invariant of X vanishes, that is, $SW_X^\circ(L) = 0$ for any characteristic line bundle L on X .*

Proof: Proofs of (i) and (ii) are exactly the same as the proofs of case $b_2^+ > 1$ as long as the SW-invariant SW_X° is well defined, i.e. it is independent of metrics on X . Let L be a characteristic line bundle on X such that the formal dimension, $\frac{1}{4}(c_1(L)^2 - 3\sigma(X) - 2e(X))$, of the moduli space is non-negative and even. The condition $b_2^+ = 1$ and $b_2^- \leq 9$ imply that $c_1(L)^2 \geq 3\sigma(X) + 2e(X) \geq 0$. Furthermore, since X is simply connected and $c_1(L)$ is characteristic, $c_1(L) \neq 0$ (Otherwise, X has $b_2^- = 9$ and it is spin which contradicts the Rohlin's signature theorem.) Thus, for any metric g on X , the light cone lemma implies $c_1(L) \cdot [\omega_g] \neq 0$, so that the sign of $(2\pi c_1(L)) \cdot [\omega_g]$ is the same for all generic metrics. Hence the SW-invariant $SW_X^\circ(L)$ is well defined. $\square$

Next we briefly review a *rational blow-down* technique introduced by R. Fintushel and R. Stern and state related facts ([FS1] for details).

Let C_p be a smooth 4-manifold obtained by plumbing the $(p-1)$ disk bundles over the 2-sphere instructed by the following diagram

$$\begin{array}{ccccccc} -(p+2) & -2 & & & & & -2 \\ \bullet & \bullet & \cdots & \cdots & \cdots & \cdots & \bullet \\ u_{p-1} & u_{p-2} & & & & & u_1 \end{array}$$

where each vertex u_i represents a disk bundle over the 2-sphere with Euler class labelled above and an interval between vertices indicates plumbing the disk bundles corresponding to the vertices. Label the homology classes represented by the 2-spheres in C_p by $u_1, \dots, u_{p-1}$ so that the self-intersections are $u_{p-1}^2 = -(p+2)$ and $u_i^2 = -2$ for $1 \leq i \leq p-2$. Furthermore, orient the 2-spheres so that $u_i \cdot u_{i+1} = +1$. Then a configuration C_p has the following topological properties:

- (1) It is a negative definite simply connected smooth 4-manifold whose boundary is the lens space $L(p^2, 1-p)$, and the lens space $L(p^2, 1-p) = \partial C_p$ bounds a rational ball B_p with $\pi_1(B_p) \cong \mathbf{Z}_p$.
- (2) $H_2(C_p; \mathbf{Z}) \cong \bigoplus_{i=1}^{p-1} \mathbf{Z}$ has generators $\{u_i : 1 \leq i \leq p-1\}$, where each u_i can be represented by the zero-section S_i^2 of the disk bundle u_i over S^2 (We use u_i for both a generator and the corresponding disk bundle).

- (3) Let P be a plumbing matrix for C_p with respect to the basis $\{u_i : 1 \leq i \leq p-1\}$. Then the intersection form on $H^2(C_p; \mathbf{Q})$ with respect to the dual basis $\{\gamma_i : 1 \leq i \leq p-1\}$ (i.e. $\langle \gamma_i, u_j \rangle = \delta_{ij}$) is given by

$$Q := (\gamma_i \cdot \gamma_j) = P^{-1}.$$

For example, when $p = 7$, we have the following intersection form on $H^2(C_7; \mathbf{Q})$ with respect to $\{\gamma_i : 1 \leq i \leq 6\}$:

$$Q_7 = \frac{-1}{49} \begin{pmatrix} 41 & 33 & 25 & 17 & 9 & 1 \\ 33 & 66 & 50 & 34 & 18 & 2 \\ 25 & 50 & 75 & 51 & 27 & 3 \\ 17 & 34 & 51 & 68 & 36 & 4 \\ 9 & 18 & 27 & 36 & 45 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix}$$

Definition Suppose X is a smooth 4-manifold which contains a configuration C_p . Then we construct a new smooth 4-manifold X_p , called the *rational blow-down* of X , by replacing C_p with a rational ball B_p . We call this a *rational blow-down* technique. Note that this process is well-defined, that is, a new smooth 4-manifold X_p is uniquely constructed (up to diffeomorphism) from X because each diffeomorphism of $\partial B_p = L(p^2, 1-p)$ extends over the rational ball B_p . Furthermore, M. Symington proved that a rational blow-down manifold X_p admits a symplectic structure in some cases.

Theorem 2.4 ([Sy]). *Suppose X is a symplectic 4-manifold containing a configuration C_p with a symplectic 2-form ω . If all 2-spheres u_i in C_p are symplectically embedded and intersect positively, then the rational blow-down manifold $X_p = X_0 \cup_{L(p^2, 1-p)} B_p$ admits a symplectic 2-form ω_p such that $(X_0, \omega_p|_{X_0})$ is symplectomorphic to $(X_0, \omega|_{X_0})$.*

Remark 1. Suppose $X = X_0 \cup_{L(p^2, 1-p)} C_p$ is a symplectic 4-manifold with a canonical class K and a compatible symplectic 2-form ω . In the case when $X_p = X_0 \cup_{L(p^2, 1-p)} B_p$ admits a symplectic structure as in the Theorem 2.4 above, let ω_p be the induced symplectic 2-form on X_p such that $\psi_p : (X_0, \omega_p|_{X_0}) \rightarrow (X_0, \omega|_{X_0})$ is a symplectomorphism. We also let K_p be the canonical class on X_p which is induced from the symplectic 2-form ω_p on X_p . Then, since $H^1(L(p^2, 1-p); \mathbf{Q}) = H^2(L(p^2, 1-p); \mathbf{Q}) = 0$, if we decompose K and ω as

$$K = K|_{X_0} + K|_{C_p} \quad \text{and} \quad [\omega] = [\omega|_{X_0}] + [\omega|_{C_p}]$$

with $K|_{X_0}, [\omega|_{X_0}] \in H^2(X_0; \mathbf{Q})$ and $K|_{C_p}, [\omega|_{C_p}] \in H^2(C_p; \mathbf{Q})$, we can also decompose K_p and ω_p as

$$K_p = K_p|_{X_0} + K_p|_{B_p} \quad \text{and} \quad [\omega_p] = [\omega_p|_{X_0}] + [\omega_p|_{B_p}].$$

Lemma 2.1. *Under the same hypothesis on (X, K, ω) and (X_p, K_p, ω_p) as above, we have*

$$K_p \cdot [\omega_p] = K \cdot [\omega] - K|_{C_p} \cdot [\omega|_{C_p}].$$


Proof: Since $K_p|_{B_p}$ and $[\omega_p|_{B_p}]$ are zero elements in $H^2(B_p; \mathbf{Q})$, we have

$$\begin{aligned} K_p \cdot [\omega_p] &= K_p|_{X_0} \cdot [\omega_p|_{X_0}] = \psi_p^*(K|_{X_0}) \cdot \psi_p^*([\omega|_{X_0}]) \\ &= K|_{X_0} \cdot [\omega|_{X_0}] = K \cdot [\omega] - K|_{C_p} \cdot [\omega|_{C_p}]. \quad \square \end{aligned}$$

3. A MAIN CONSTRUCTION

In this section we construct a new family of simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$ which are homeomorphic, but not diffeomorphic, to rational surfaces by using rational blow-down technique introduced by R. Fintushel and R. Stern in [FS1].

Let us start with analyzing a simply connected rational surface $E(1) = \mathbf{CP}^2 \# 9\overline{\mathbf{CP}}^2$. There are several ways to describe $E(1)$. One way to construct $E(1)$ is to take two generic cubic curves in $\mathbf{CP}^2$ which intersect each other at 9 points and then blow up 9 times at these points in $\mathbf{CP}^2$. This viewpoint makes us to see $E(1) = \mathbf{CP}^2 \# 9\overline{\mathbf{CP}}^2$ as a Lefschetz fibration over $\mathbf{CP}^1$ whose generic fiber is an elliptic curve, say f , and which also has 6 singular cusp fibers (or, equivalently, 12 singular fishtail fibers). Since 4 singular cusp fibers in $E(1)$ can be deformed to an $\widetilde{E}_6$ -singular fiber, $E(1)$ can also be described as an elliptic fibration over $\mathbf{CP}^1$ with 3 singular fibers, one $\widetilde{E}_6$ -fiber and two cusp fibers. Note that a neighborhood of the $\widetilde{E}_6$ -fiber in $E(1)$ is a smooth 4-manifold obtained by plumbing disk bundles over the holomorphically embedded 2-spheres $S_i (1 \leq i \leq 7)$ of square -2 instructed by the Dynkin diagram of $\widetilde{E}_6$ ([HKK] for details).

FIGURE 1. $\widetilde{E}_6$ -singular fiber

Lemma 3.1 ([A], [F]). *The second (co)homology classes $[S_i]$ ($1 \leq i \leq 7$) of the 2-spheres S_i embedded in $\widetilde{E}_6$ can be represented by $[S_1] = e_4 - e_7$, $[S_2] = e_1 - e_4$, $[S_3] = h - e_1 - e_2 - e_3$, $[S_4] = e_2 - e_5$, $[S_5] = e_5 - e_9$, $[S_6] = e_3 - e_6$ and $[S_7] = e_6 - e_8$, where h denotes a generator of $H_2(\mathbf{CP}^2; \mathbf{Z})$ and each e_i denotes the (co)homology class represented by the i^{th} exceptional curve in $\overline{\mathbf{CP}}^2 \subset E(1) = \mathbf{CP}^2 \# 9\overline{\mathbf{CP}}^2$.*

Proof: Note that $E(1)$ can be constructed as follows: First choose a generic cubic curve C (represented homologically by $3h$) which intersects with a line (represented by h) at 3 points in $\mathbf{CP}^2$. And then blow up at these 3 points, so that we get an embedded 2-sphere S_3 , represented by $h - e_1 - e_2 - e_3$, of multiplicity 3 and of square -2 in $\mathbf{CP}^2 \# 3\overline{\mathbf{CP}}^2$. Again blow up 3 times at the intersection points between the curve $C - e_1 - e_2 - e_3$ and 3 exceptional curves e_1, e_2 and e_3 respectively, so that we get embedded 2-spheres $S_2 = e_1 - e_4, S_4 = e_2 - e_5$ and $S_6 = e_3 - e_6$ of multiplicities 2 and of squares -2 in $\mathbf{CP}^2 \# 6\overline{\mathbf{CP}}^2$. Finally blow up 3 times at the intersection points between the curve $C - e_1 - e_2 - \dots - e_6$ and 3 new exceptional curves e_4, e_5 and e_6 respectively, so that we get again embedded 2-spheres $S_1 = e_4 - e_7, S_5 = e_5 - e_9$ and $S_7 = e_6 - e_8$ of multiplicities 1 and of squares -2 in $\mathbf{CP}^2 \# 9\overline{\mathbf{CP}}^2$. Then the embedded 2-spheres $\{S_1 \dots S_7\}$ consists of $\widetilde{E}_6$ -singular fiber in $E(1)$. $\square$

Note that the standard canonical class $K_{E(1)} \in H^2(E(1); \mathbf{Z})$ of $E(1)$ is represented by $K_{E(1)} = -3h + (e_1 + \cdots + e_9) = -[f]$. Furthermore, there is a relation between the canonical class and a compatible symplectic 2-form on a non-minimal rational surface which will play an important role in the proof of our main results. Explicitly, we have

Lemma 3.2. *For each integer $k \geq 1$, there exists a symplectic 2-form ω on $E(1) \# k \overline{\mathbf{CP}}^2$ which is compatible with the standard canonical class $K_{E(1) \# k \overline{\mathbf{CP}}^2} = -3h + (e_1 + \cdots + e_{9+k})$ such that its cohomology class $[\omega]$ can be represented by $ah - (b_1 e_1 + \cdots + b_{9+k} e_{9+k})$ for some rational numbers $a, b_1, \dots, b_{9+k}$ satisfying $a \geq b_1 \geq b_2 \geq \dots \geq b_{9+k} \geq 0$ and $3a > b_1 + \cdots + b_{9+k}$.*

Proof : Since $E(1) \# k \overline{\mathbf{CP}}^2$ is a rational surface, there exists a symplectic 2-form ω on $E(1) \# k \overline{\mathbf{CP}}^2$ which is compatible with the standard canonical class $K_{E(1) \# k \overline{\mathbf{CP}}^2} = -3h + (e_1 + \cdots + e_{9+k})$ satisfying $K_{E(1) \# k \overline{\mathbf{CP}}^2} \cdot [\omega] < 0$ (refer to Corollary 1.4 in [MS]). Furthermore, Lemma 4.7 in [LL2] guarantees that the cohomology class $[\omega]$ of the symplectic 2-form ω can be represented by $ah - (b_1 e_1 + \cdots + b_{9+k} e_{9+k})$ for some rational numbers $a, b_1, \dots, b_{9+k}$ satisfying $a \geq b_1 \geq b_2 \geq \dots \geq b_{9+k} \geq 0$. The inequality $3a > b_1 + \cdots + b_{9+k}$ follows from the fact that $K_{E(1) \# k \overline{\mathbf{CP}}^2} \cdot [\omega] < 0$. $\square$


In fact, T. Li and A. Liu obtained many results regarding symplectic structures and canonical classes on symplectic 4-manifolds with $b_2^+ = 1$. For example, they proved

Lemma 3.3 ([LL1]). *There is a unique symplectic structure on $\mathbf{CP}^2 \# k \overline{\mathbf{CP}}^2$ for $2 \leq k \leq 9$ up to diffeomorphisms and deformation. For $k \geq 10$, the symplectic structure is still unique for the standard canonical class. In particular, $\mathbf{CP}^2 \# k \overline{\mathbf{CP}}^2$ ($2 \leq k \leq 9$) does not admit a symplectic 2-form ω for which $c_1(K) \cdot [\omega] > 0$.*

Proposition 3.1. *There exists a configuration C_7 in a rational surface $E(1) \# 4 \overline{\mathbf{CP}}^2$ such that all 2-spheres u_i lying in C_7 are symplectically embedded.*

Proof : Note that $E(1)$ can be viewed as an elliptic fibration with an $\widetilde{E}_6$ -singular fiber and 2 cusp singular fibers (equivalently, 4 singular fishtail fibers). Since the homology class $[f]$ of the elliptic fiber f in $E(1)$ can be represented by an immersed 2-sphere with one positive double point (equivalently, a fishtail fiber) and since $E(1)$ contains at least 4 such immersed 2-spheres, we blow up 4 times at these double points so that there exist embedded 2-spheres, $f - 2e_{10}, \dots, f - 2e_{13}$, in $E(1) \# 4 \overline{\mathbf{CP}}^2$ which intersect a section e_9 of $E(1)$ positively at points, say $p_1, \dots, p_4$, respectively. And then, resolving symplectically the intersection points $p_1, \dots, p_4$ between $f - 2e_{10}, \dots, f - 2e_{13}$ and e_9 , we have a symplectically embedded 2-sphere, denoted by S , in $E(1) \# 4 \overline{\mathbf{CP}}^2$ which represents a homology class $(f - 2e_{10}) + \cdots + (f - 2e_{13}) + e_9 = 4f + e_9 - 2(e_{10} + \cdots + e_{13})$ with square -9 . Now, using a linear plumbing manifold consisting of 5 disk bundles $\{S_1, S_2, \dots, S_5\}$ lying in a neighborhood of an $\widetilde{E}_6$ -singular fiber (Figure 1), we obtain a configuration $C_7 \subset E(1) \# 4 \overline{\mathbf{CP}}^2$ by setting $u_1 = S_1, u_2 = S_2, \dots, u_5 = S_5$ and $u_6 = S$ (Figure 2). Note that all 2-spheres u_i lying in the configuration C_7 are symplectically embedded. $\square$

Remark 2. Note that there are other candidates for a configuration C_7 in $E(1) \# 4 \overline{\mathbf{CP}}^2$ by choosing a different linear plumbing manifold lying in a neighborhood of an $\widetilde{E}_6$ -singular

FIGURE 2. $C_7 \subset E(1)\sharp 4\overline{\mathbf{CP}}^2$

fiber. For example, one may choose a linear plumbing manifold consisting of 5 disk bundles $\{S_1, S_2, S_3, S_6, S_7\}$ in Figure 1 to get a configuration C_7 .

Theorem 3.1. *There exists a simply connected symplectic 4-manifold with $b_2^+ = 1$ and $c_1^2 = 2$ which is homeomorphic, but not diffeomorphic, to $\mathbf{CP}^2\sharp 7\overline{\mathbf{CP}}^2$.*

Proof : Construction - By Proposition 3.1 above, we have a symplectically embedded configuration C_7 in a rational surface $X := E(1)\sharp 4\overline{\mathbf{CP}}^2$. Hence we get a new smooth 4-manifold, denoted by $X_7 = X_0 \cup_{L(49, -6)} B_7$, by rationally blowing down along the configuration C_7 in $X = X_0 \cup_{L(49, -6)} C_7$. Furthermore, Theorem 2.4 guarantees the existence of a symplectic structure on X_7 .

Properties of X_7 - Since a circle representing a generator of $\pi_1(L(49, -6))$ bounds a disk which is a hemisphere of S_6 lying in $\widetilde{E}_6$ -singular fiber, $\pi_1(X_0) = 1$. Hence the simple connectivity of X_7 follows from Van-Kampen's theorem. Furthermore, it satisfies $c_1^2(X_7) = c_1^2(X) + 6 = 2$, so that it is homeomorphic to $\mathbf{CP}^2\sharp 7\overline{\mathbf{CP}}^2$ due to M. Freedman's classification theorem of simply connected closed topological 4-manifolds. It only remains to show that X_7 is not diffeomorphic to $\mathbf{CP}^2\sharp 7\overline{\mathbf{CP}}^2$. For this, we first claim that the canonical class K_7 on X_7 and the corresponding symplectic 2-form ω_7 on X_7 satisfy $K_7 \cdot [\omega_7] > 0$. Then, since $b_2^-(X_7) \leq 9$ and $(-K_7) \cdot [\omega_7] < 0$, the Seiberg-Witten invariant $SW_{X_7}^\circ$ is well defined and we have $SW_{X_7}^\circ(-K_7) = SW_{X_7}^{\circ,-}(-K_7) = SW_{X_7}^-(-K_7) = \pm 1$, where the last equality follows from Theorem 2.2. Note that the non-triviality of $SW_{X_7}^\circ$ means that X_7 does not admit a metric of positive scalar curvature, equivalently, it is not diffeomorphic to a rational surface. Hence we are done.

Proof of $K_7 \cdot [\omega_7] > 0$ - Note that the canonical class K of $X = E(1)\sharp 4\overline{\mathbf{CP}}^2$ is represented by $K = -3h + (e_1 + \cdots + e_{13}) = -[f] + (e_{10} + \cdots + e_{13})$ and, by modifying B. Li and T. Li's symplectic cone argument in [LL2], we may assume that the cohomology class $[\omega]$ of a symplectic 2-form ω on X compatible with a canonical class K can be represented by $ah - (b_1e_1 + \cdots + b_{13}e_{13})$ for some rational numbers $a, b_1, \dots, b_{13}$ satisfying $a \geq b_1 \geq b_2 \geq \cdots \geq b_{13} \geq 0$ and $3a > b_1 + \cdots + b_{13}$ (refer to Lemma 3.2). Now, remembering that $u_6 = S = 4f + e_9 - 2(e_{10} + \cdots + e_{13}) = 12h + e_9 - 4(e_1 + \cdots + e_9) - 2(e_{10} + \cdots + e_{13})$ in Proposition 3.1 and the canonical class K does not intersect with holomorphic 2-spheres u_i ($1 \leq i \leq 5$) of square -2 , let us express two cohomology classes $K|_{C_7}$ and $[\omega]|_{C_7}$ using a dual basis $\{\gamma_i : 1 \leq i \leq 6\}$ (i.e. $\langle \gamma_i, u_j \rangle = \delta_{ij}$) for $H^2(C_7; \mathbf{Q})$:

$$\begin{aligned}
K|_{C_7} &= (K \cdot u_1)\gamma_1 + (K \cdot u_2)\gamma_2 + \cdots + (K \cdot u_6)\gamma_6 \\
&= 7\gamma_6 \text{ and} \\
[\omega]|_{C_7} &= ([\omega] \cdot u_1)\gamma_1 + ([\omega] \cdot u_2)\gamma_2 + \cdots + ([\omega] \cdot u_5)\gamma_5 + ([\omega] \cdot u_6)\gamma_6 \\
&= (b_4 - b_7)\gamma_1 + (b_1 - b_4)\gamma_2 + (a - b_1 - b_2 - b_3)\gamma_3 + (b_2 - b_5)\gamma_4 \\
&\quad + (b_5 - b_9)\gamma_5 + \{12a - 4(b_1 + \cdots + b_9) - 2(b_{10} + \cdots + b_{13}) + b_9\}\gamma_6.
\end{aligned}$$

Then, using the intersection form Q_7 on $H^2(C_7; \mathbf{Q})$, we have

$$\begin{aligned} K|_{C_7} \cdot [\omega|_{C_7}] &= \frac{-1}{7} \{ (b_4 - b_7) + 2(b_1 - b_4) + 3(a - b_1 - b_2 - b_3) + 4(b_2 - b_5) \\ &\quad + 5(b_5 - b_9) + 6(12a - 4(b_1 + \cdots + b_9) - 2(b_{10} + \cdots + b_{13}) + b_9) \} \\ &= \frac{-1}{7} \{ 75a - 25b_1 - 23b_2 - 27b_3 - 25b_4 - 23b_5 - 24b_6 - 25b_7 - 24b_8 \\ &\quad - 23b_9 - 12(b_{10} + \cdots + b_{13}) \}. \end{aligned}$$

Hence, by Lemma 2.1 and Lemma 3.2, we have

$$\begin{aligned} K_7 \cdot [\omega_7] &= K \cdot [\omega] - K|_{C_7} \cdot [\omega|_{C_7}] \\ &= \{ -3a + (b_1 + \cdots + b_{13}) \} - K|_{C_7} \cdot [\omega|_{C_7}] \\ &= \frac{1}{7} \{ 54a - 18b_1 - 16b_2 - 20b_3 - 18b_4 - 16b_5 - 17b_6 - 18b_7 - 17b_8 \\ &\quad - 16b_9 - 5(b_{10} + \cdots + b_{13}) \} \\ &> \frac{1}{7} \{ 2b_2 - 2b_3 + 2b_5 + b_6 + b_8 + 2b_9 + 13(b_{10} + \cdots + b_{13}) \} \\ &\geq 0. \quad \square \end{aligned}$$

Remark 3. Since the canonical class K_7 induced from a symplectic structure ω_7 on the symplectic 4-manifold X_7 constructed in the proof of Theorem 3.1 above satisfies $K_7 \cdot [\omega_7] > 0$, one can also conclude directly from Lemma 3.3 that X_7 is not diffeomorphic to a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$.

Remark 4. Similarly, using various different configurations C_7 lying in $E(1) \# 4\overline{\mathbf{CP}}^2$ (refer to Remark 2) and the same technique as in the proof of Theorem 3.1 above, we can construct a family of simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$ which are all homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$. But we do not know whether all these symplectic 4-manifolds are mutually diffeomorphic to each other.

Remark 5. There are still some important questions to be solved regarding on the symplectic 4-manifold X_7 . For example, though it is likely to be minimal, it is not easy to determine whether X_7 is minimal. Furthermore, it is a very intriguing question whether X_7 admits a complex structure.

Remark 6. Theorem 3.1 above enables us to confirm that a rational surface $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$ admits an exotic smooth structure.

4. SYMPLECTIC 4-MANIFOLDS WITH $b_2^+ = 1$ AND $c_1^2 = 1$

As we mentioned in the Introduction, the only known simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 1$ are complex surfaces such as a rational surface $\mathbf{CP}^2 \# 8\overline{\mathbf{CP}}^2$ and Barlow surfaces. In this section, using the same technique as in the proof of Theorem 3.1 above, we construct simply connected symplectic 4-manifolds which are homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2 \# 8\overline{\mathbf{CP}}^2$.

Proposition 4.1. *There exists a configuration C_5 in a rational surface $E(1)\#3\overline{\mathbf{CP}}^2$ such that all 2-spheres u_i lying in C_5 are symplectically embedded.*

Proof : As in the proof of Proposition 3.1, we blow up 3 times at the double points of singular fishtail fibers so that there exist embedded 2-spheres, $f - 2e_{10}, \dots, f - 2e_{12}$, in $E(1)\#3\overline{\mathbf{CP}}^2$ which intersect a section e_2 of $E(1)$ positively at points, say $p_1, \dots, p_3$, respectively. And then, resolving symplectically the intersection points $p_1, \dots, p_3$ between $f - 2e_{10}, \dots, f - 2e_{12}$ and e_2 , we have a symplectically embedded 2-sphere, denoted by S , in $E(1)\#3\overline{\mathbf{CP}}^2$ which represents a homology class $(f - 2e_{10}) + \dots + (f - 2e_{12}) + e_2 = 3f + e_2 - 2(e_{10} + \dots + e_{12})$ with square -7 . Now, using a linear plumbing manifold consisting of 3 disk bundles $\{S_1, S_2, S_3\}$ lying in a neighborhood of an $\widetilde{E}_6$ -singular fiber (Figure 1), we obtain a configuration $C_5 \subset E(1)\#3\overline{\mathbf{CP}}^2$ by setting $u_1 = S_1, u_2 = S_2, u_3 = S_3$ and $u_4 = S$. $\square$

Theorem 4.1. *There exists a simply connected symplectic 4-manifold with $b_2^+ = 1$ and $c_1^2 = 1$ which is homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$.*

Proof : By Proposition 4.1 above, we have a symplectically embedded configuration C_5 in a rational surface $X := E(1)\#3\overline{\mathbf{CP}}^2$. Hence we get a new smooth 4-manifold, denoted by $X_5 = X_0 \cup_{L(25, -4)} B_5$, by rationally blowing down along the configuration C_5 in $X = X_0 \cup_{L(25, -4)} C_5$. Then the rest of proof is exactly the same as the proof of Theorem 3.1 above except a computation of $K_5 \cdot [\omega_5] > 0$, which is the following:

In this case, we have $K = -3h + (e_1 + \dots + e_{12})$ and $[\omega] = ah - (b_1e_1 + \dots + b_{12}e_{12})$ for some rational numbers $a, b_1, \dots, b_{12}$ satisfying $a \geq b_1 \geq b_2 \geq \dots \geq b_{12} \geq 0$ and $3a > b_1 + \dots + b_{12}$. Now, using $u_4 = S = 3f + e_2 - 2(e_{10} + \dots + e_{12}) = 9h + e_2 - 3(e_1 + \dots + e_9) - 2(e_{10} + \dots + e_{12})$, we have

$$\begin{aligned} K|_{C_5} &= (K \cdot u_1)\gamma_1 + \dots + (K \cdot u_4)\gamma_4 = 5\gamma_4 \quad \text{and} \\ \omega|_{C_5} &= ([\omega] \cdot u_1)\gamma_1 + \dots + ([\omega] \cdot u_4)\gamma_4 \\ &= (b_4 - b_7)\gamma_1 + (b_1 - b_4)\gamma_2 + (a - b_1 - b_2 - b_3)\gamma_3 \\ &\quad + \{9a - 3(b_1 + \dots + b_9) - 2(b_{10} + \dots + b_{12}) + b_2\}\gamma_4. \end{aligned}$$

Hence, by Lemma 2.1 and Lemma 3.2, we get

$$\begin{aligned} K|_{C_5} \cdot [\omega|_{C_5}] &= \frac{-1}{5} \{ (b_4 - b_7) + 2(b_1 - b_4) + 3(a - b_1 - b_2 - b_3) \\ &\quad + 4(9a - 3(b_1 + \dots + b_9) - 2(b_{10} + \dots + b_{12}) + b_2) \} \\ &= \frac{-1}{5} \{ 39a - 13b_1 - 11b_2 - 15b_3 - 13b_4 - 12(b_5 + b_6) - 13b_7 \\ &\quad - 12(b_8 + b_9) - 8(b_{10} + \dots + b_{12}) \}, \\ K_5 \cdot [\omega_5] &= K \cdot [\omega] - K|_{C_5} \cdot [\omega|_{C_5}] \\ &= \frac{1}{5} \{ 24a - 8b_1 - 6b_2 - 10b_3 - 8b_4 - 7(b_5 + b_6) - 8b_7 \\ &\quad - 7(b_8 + b_9) - 3(b_{10} + \dots + b_{12}) \} \\ &> \frac{1}{5} \{ 2b_2 - 2b_3 + b_5 + b_6 + b_8 + b_9 + 5(b_{10} + \dots + b_{12}) \} \\ &\geq 0. \quad \square \end{aligned}$$

Remark 7. Similarly, using various different configurations C_5 lying in $E(1)\#3\overline{\mathbf{CP}}^2$, we can also construct a family of simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 1$ which are all homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$. But we do not know whether one of these symplectic 4-manifolds is diffeomorphic to a Barlow surface.

Finally, we also investigate exotic smooth structures on a rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$. As mentioned in the Introduction, Barlow surface, denoted by B , is homeomorphic, but not diffeomorphic, to a rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$. Furthermore, whereas $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$ admits an Einstein metric with positive scalar curvature, B admits an Einstein metric with negative scalar curvature. Now, since a symplectic 4-manifold X_7 constructed in Theorem 3.1 has a nontrivial SW-basic class K_7 obtained by a small generic perturbation, so that its blow-up manifold $X_7\#\overline{\mathbf{CP}}^2$ has at least two (up to sign) SW-basic classes, $K_7 + E$ and $K_7 - E$ (E is an exceptional curve obtained by a blowing up), we conclude that the blow-up manifold $X_7\#\overline{\mathbf{CP}}^2$ is also homeomorphic, but not diffeomorphic, to both B and $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$. Furthermore, D. Kotschick pointed out that $X_7\#\overline{\mathbf{CP}}^2$ does not admit an Einstein metric, which can be deduced from his result regarding on the existence of Einstein metrics (Theorem 4.3 below for details). Hence we get the following useful information about exotic smooth structures on a rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$. Before we state our final result, we first quote a theorem of D. Kotschick:

Theorem 4.2 ([K2]). *Let X be a smooth 4-manifold with a monopole class c . If X admits an Einstein metric, then the maximal number k of copies of $\overline{\mathbf{CP}}^2$ that can be split off smoothly is bounded by*

$$k \leq \frac{1}{2}\{2e(X) + 3\sigma(X) - 8d\}$$

where d is the dimension of moduli space of the solutions of Seiberg-Witten equations corresponding to a monopole class c .

Theorem 4.3. *A rational surface $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$ admits at least three distinct smooth structures - an Einstein metric with positive scalar curvature, an Einstein metric with negative scalar curvature and no Einstein metric.*

Proof : Since $X_7\#\overline{\mathbf{CP}}^2$, B and $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$ are not mutually diffeomorphic to each other, and since B and $\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$ have desired Einstein metrics, it only suffices to show that $X_7\#\overline{\mathbf{CP}}^2$ does not admit an Einstein metric. For this, suppose that a symplectic 4-manifold $X_7\#\overline{\mathbf{CP}}^2$ has an Einstein metric and apply Theorem 4.2 above to $X_7\#\overline{\mathbf{CP}}^2$. Then we have $1 \leq k \leq \frac{1}{2}\{2e(X) + 3\sigma(X) - 8d\} = \frac{1}{2}$, which is a contradiction. $\square$

REFERENCES

- [A] D. Auroux, Personal communication
- [F] R. Fintushel, Personal communication
- [FS1] R. Fintushel and R. Stern, Rational blowdowns of smooth 4-manifolds, Jour. Diff. Geom. **46** (1997), 181–235
- [FS2] R. Fintushel and R. Stern, Knots, links and 4-manifolds, Invent. Math. **134** (1998), 363–400

- [G] R. Gompf, A new construction of symplectic manifolds, *Annals of Math.* **142** (1995), 527–595
- [HKK] J. Harer, A. Kas and R. Kirby, Handlebody decompositions of complex surfaces, *Memoirs of the AMS* **62** (1986), AMS
- [Ka] W. Kazez, *Geometric Topology - Part 2, Studies in Advanced Mathematics* **2.2** (1997), AMS/IP
- [K1] D. Kotschick, On manifolds homeomorphic to $\mathbf{CP}^2 \# 8\overline{\mathbf{CP}}^2$, *Invent. Math.* **95** (1989), 591–600
- [K2] D. Kotschick, Monopole classes and Einstein metrics, preprint (2003), arXiv:math.DG/0306022
- [LL1] T. Li and A. Liu, Symplectic structures on ruled surfaces and a generalized adjunction formula, *Math. Res. Letters* **2** (1995), 453–471
- [LL2] T. Li and A. Liu, Uniqueness of symplectic canonical class, surface cone and symplectic cone of 4-manifolds with $b^+ = 1$, *Jour. Diff. Geom.* **58** (2001), 331–370
- [M] J. Morgan, *The Seiberg-Witten equations and applications to the topology of smooth four-manifolds*, *Math. Notes* **44**, Princeton University Press, 1996
- [MS] D. McDuff and D. Salamon, A survey of symplectic 4-manifolds with $b_2^+ = 1$, *Turkish Jour. Math.* **20** (1996), 47–61
- [Sy] M. Symington, Symplectic rational blowdowns, *Jour. Diff. Geom.* **50** (1998), 505–518
- [Sz] Z. Szabó, Exotic 4-manifolds with $b_2^+ = 1$, *Math. Res. Letters* **3** (1996), 731–741
- [T] C.H. Taubes, The Seiberg-Witten invariants and symplectic forms, *Math. Res. Letters* **1** (1994), 809–822

ADDRESS 1: DEPARTMENT OF MATHEMATICS, KONKUK UNIVERSITY, 1 HWAYANG-DONG, KWANGJIN-GU, SEOUL 143-701, KOREA

ADDRESS 2: DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI48824, USA

E-mail address: jipark@konkuk.ac.kr, jipark@math.msu.edu